



A Study to Assess the Frequency of Sleep Disturbances in Down Syndrome Children

¹Dr. Vivechana Kaniyala, ²Dr. Anil Kumar H, ³Dr. Mallesh Kariyappa

¹PG Scholar, Dept. of Pediatrics, Bangalore Medical College and Research Institute, Bangalore, India

²Assistant Professor, Dept. of Pediatrics, Bangalore Medical College and Research Institute, Bangalore, India

³Professor and HOD, Dept. of Pediatrics, Bangalore Medical College and Research Institute, Bangalore, India

Citation of this Article: Dr. Vivechana Kaniyala, Dr. Anil Kumar H, Dr. Mallesh Kariyappa, “A Study to Assess the Frequency of Sleep Disturbances in Down Syndrome Children,” IJMSAR – May – 2023, Vol. – 6, Issue - 3, Page No. 24-27.

Copyright: © 2023, Dr. Vivechana Kaniyala, et al. This is an open access journal and article distributed under the terms of the creative common attribution noncommercial License. This allows others to remix, tweak, and build upon the work non commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

Corresponding Author: Dr. Vivechana Kaniyala, PG Scholar, Dept. of Pediatrics, Bangalore Medical College and Research Institute, Bangalore, India

Type of Publication: Original Research Article

Conflicts of Interest: Nil

ABSTRACT

Introduction: Down syndrome or trisomy 21 is the commonest chromosomal abnormality. Sleep problems are very common in these children. According to National Institute of Health 76% of children with Down syndrome experience sleep disturbances. (1,2) This study is conducted to know the frequency of sleep disturbances in children with Down syndrome at our center.

Method

A cross sectional study was conducted, data was collected regarding sleep habits of 43 Down syndrome children attending District Early Intervention Centre (DEIC) at Vanivilas Childrens Hospital, Bangalore. Sleep disturbances in these children will be assessed using Sleep Disturbances Scale for Children (SDSC).

Assessment tool were evaluated, variables assessed using chi-square test.

Results

There is significant sleep disturbance in children with Down syndrome. 30.8% preschool children and 17.6% of children above 6 years of age were found to have significant sleep problems. Disorders of initiation of sleep and Excessive somnolence were the most common sleep disorders in our study.

Conclusion

Parents report a significant sleep disturbance in both preschool and school aged children with Down syndrome. Pediatricians should routinely enquire about sleep habits in such children.

Keywords

Down syndrome, Sleep, Sleep Disturbances Scale for Children

INTRODUCTION

Down syndrome or trisomy 21 is the commonest chromosomal abnormality occurring in humans. They present with wide range of clinical signs and symptoms like developmental and intellectual disabilities, neurological features, cardiac defects etc.(1)

Sleep problems are commonly seen in children with Down syndrome. According to National Institute of Health (2)76% of children with Down syndrome experience difficulty with sleep onset and ability to stay asleep. In childhood, behavioral sleep problems are more common whereas in adults sleep related breathing problems occur frequently. Sleep disorders have a negative impact on general and mental health, behavioral and cognitive functions. Increase in hyperactivity, aggression, irritability is seen in such children. It also contributes to family burden and more social isolation. (2,3,4,5,6,7) Sleep disturbances can accelerate the progression of dementia and Alzheimer's in people with Down syndrome. Diagnosis and treatment of sleep disorders is thus essential to prevent further consequences and comorbidities. Improving overall sleep quality is the key in optimization of physical and cognitive functioning of children with this syndrome.(8)

This study is thus conducted to know the prevalence of sleep disturbances in children with Down syndrome at our hospital.

METHODOLOGY

Study design: We conducted a cross sectional Study on children with Down Syndrome attending our hospital.

Study period: Study was done during April 2023.

Place of study: Dept. of Pediatrics, Vani Vilas Children Hospital, BMCRI

Study population: Children aged between 1 to 18 years diagnosed with Down syndrome attending our hospital.

Data: Data was collected regarding the sleep habits of children with Down syndrome using Sleep Disturbances Scale for Children (SDSC).

Statistical analysis: Assessment tool were evaluated, variables were assessed using chi-square test. Association between categorical variables were assessed using chi-square test. There are 43 participants in this study (n=43). Shapiro-Wilk test was used to determine the normality of the data. CHI-SQUARE test was used to check the association between categorical variables. Continuous variables are expressed in terms of Mean, Std deviation (SD), Median, Interquartile range (IQR). Categorical variables are expressed in frequency (n) and percentage (%)

RESULTS

43 children with Down syndrome participated in the study. Participants were divided into two groups of 0-6 years and 6 -18years based on physiological sleep patterns. Of the 43 participants, 26 (60.5%) were below the age of 6 years and 17 (39.5%) were above 6 years of age. There were 20 (46.5%) girls and 23(53.5%) boys in the study. SDSC questionnaire was used to assess the sleep habits in these children. A score of 39 and above is considered to be significant. Particular sleep problems were notable: 26 children (60.3%) needed an average of >30 minutes time to fall asleep. A score of >39 was seen in 11(48.4%) children which is of great concern. These disturbances were noted in both the age groups. Children aged > 6 years

of age had marginally higher disorders of initiating and maintaining sleep (mean of 12.04 and SD of 3.21) as compared to children aged below 6 years (mean of 10.41 with SD of 2.7). Excessive somnolence was seen more in children aged above 6 years (mean of 9.31 with SD of 1.64) than children below 6 years (mean of 8.94 with SD of 1.52).

	>6 years		< 6 years	
	Mean	SD	Mean	SD
DIMS	12.04	3.21	10.41	2.72
SBD	4.50	0.65	4.41	0.51
DA	3.34	0.58	3.31	0.77
SWTD	6.54	0.91	7.00	1.60
DOES	9.31	1.64	8.94	1.52
SHY	2.42	0.64	2.35	0.50
TOTAL	38.38	3.87	37.06	3.31

DISCUSSION

Sleep is an essential component of overall health. Sleep is also necessary for overall mental and physical health as poor sleep habits can contribute to major and chronic health problems. Studies have shown that children with Down syndrome experience significant sleep problems. In our study we used a parent report questionnaire to assess these problems. SDSC is a 27-item questionnaire which assesses the sleep habit of the child over the last 6 months. Questions specific to initiation and maintenance of sleep, breathing disorders, disorders of arousal, sleep wake transition disorders excessive somnolence and hyperhidrosis are asked. Elisa Fuca et al conducted a retrospective cross-sectional study of sleep and behavioral problems in preschool children with Down syndrome at Pediatric Unit of Bambino Gesù Children's Hospital, Italy between June to November 2021. 71 children aged 3 to 5.1 years were studied using SDSC scale and Child Behavior Checklist. Descriptive statistics

were used to analyze demographical characteristics. 15.4% exhibited significant sleep problems in this study and 14.1% showed behavioral problems (7). In our study, sleep difficulty was reported in 48.4% of the children. Disorders of initiation and maintenance of sleep and excessive somnolence were the most frequently seen. Other sleep disorders in children with Down syndrome include breathing disorders such as obstructive sleep apneas, hyperhidrosis in sleep and sleep wake transition disorders such as sleepwalking, sleep talking, bruxism etc..

CONCLUSION

Sleep disorders are commonly seen in children with Down syndrome and have a significant impact on the quality of life in these children. Early identification of such problems is thus necessary. Pediatricians should routinely enquire about sleep habits in such children and intervene early to avoid long term complications.

REFERENCES

1. Rafii MS, Kleschevnikov AM, Sawa M, Mobley WC. Down syndrome. *Handb Clin Neurol*. 2019;167:321-336. doi: 10.1016/B978-0-12-804766-8.00017-0. PMID: 31753140.
2. Zipperer, H. (2019) Common sleep challenges in children with down syndrome (and what to do about them) -The Down Syndrome Association of Greater Charlotte - DSA Greater Charlotte.
3. Bassell JL, Phan H, Leu R, Kronk R, Visootsak J. Sleep profiles in children with Down syndrome. *Am J Med Genet A*. 2015 Aug;167A(8):1830-5. doi: 10.1002/ajmg.a.37096. Epub 2015 Jun 23. PMID: 26105013; PMCID: PMC4515186
4. Carter M, McCaughey E, Annaz D, Hill CM. Sleep problems in a Down syndrome population. *Arch Dis Child*. 2009 Apr;94(4):308-10. doi:

- 10.1136/adc.2008.146845. Epub 2008 Sep 11.
PMID: 18786953
5. Breslin JH, Edgin JO, Bootzin RR, Goodwin JL, Nadel L. Parental report of sleep problems in Down syndrome. *J Intellect Disabil Res.* 2011 Nov;55(11):1086-91. doi: 10.1111/j.1365-2788.2011.01435.x. Epub 2011 Jul 5. PMID: 21726315.
 6. Giménez S, Altuna M, Blessing E, Osorio RM, Fortea J. Sleep Disorders in Adults with Down Syndrome. *J Clin Med.* 2021 Jul 6;10(14):3012. doi: 10.3390/jcm10143012. PMID: 34300177; PMCID: PMC8306783
 7. Fucà E, Costanzo F, Ursumando L, Celestini L, Scoppola V, Mancini S, Valentini D, Villani A, Vicari S. Sleep and behavioral problems in preschool-age children with Down syndrome. *Front Psychol.* 2022 Jul 18;13:943516. doi: 10.3389/fpsyg.2022.943516. PMID: 35923741; PMCID: PMC9342601.
 8. Behrman, R.E. et al. (2020) *Nelson Textbook of Pediatrics*. Philadelphia: Elsevier Inc.